




Paralytic Shellfish Toxins

The paralytic shellfish toxins are a group of 18 secondary metabolites deposited in bivalve mollusks by dinoflagelates. Dinoflagelates blooms are seasonal, occurring during warm months. Since it is unpredictable whether an infestation will occur, the shellfish population should be regularly monitored for toxins. Ingestion of contaminated shellfish can lead to paralytic shellfish poisoning; a life-threatening illness.



Dinoflagellate-derived Neurotoxins in Bivalve Mollusks

METHOD

Analytical Conditions

COLUMN: Reversed-phase column for PST analysis, C₈, 4.6 x 150 mm, Catalog No. 0846150
TEMPERATURE: 40 °C
FLOW RATE: 0.8 mL/min
MOBILE PHASE: A. For C1–C4 toxins:
Tetrabutylammonium phosphate (1 mM) adjusted to pH 5.8 with acetic acid
B. For GTX-1 to GTX-6, dcGTX2, and dcGTX3:
Sodium 1-heptanesulfonate (2 mM) in 10 mM Ammonium phosphate, pH 7.1
C. For STX, neoSTX, and dcSTX:
Sodium 1-heptanesulfonate (2 mM) in 10 mM Ammonium phosphate, pH 7.1 : Acetonitrile (2 : 1)

Post-column Conditions

POST-COLUMN SYSTEM: PCX5200
REAGENT 1: Periodic acid (7 mM) in 50 mM Potassium Phosphate buffer, pH 9.0
REACTOR 1: 75 °C, 2.0 mL
REAGENT 2: 0.5 M Acetic Acid
REACTOR 2: ambient, 0.1 mL
FLOW RATE: 0.4 mL/min each
DETECTION: Fluorometer
 λ_{ex} : 330 nm
 λ_{em} : 390 nm

REFERENCES:

- 1) Y. Oshima, K Sugino, T. Yasumoto
"Latest Advances in HPLC Analysis of Paralytic Shellfish Toxins in Bioactive Molecules" **10**, (1989) 319–326
- 2) J. J. Sullivan, M. M. Wekell, and L. L. Kentala,
J. Food Sci. **50**(1) (1985) 26–29.
- 3) Y. Oshima, *JOAC, INTL.* **78**(2) (1995).